

Reproduction and growth of dusky flathead (*Platycephalus fuscus*) in NSW estuaries

Charles A. Gray & Lachlan M. Barnes

Primary Industries Science & Research
Cronulla Fisheries Research Centre of Excellence
P.O. Box 21, Cronulla, NSW 2230
Australia



July 2008

NSW Department of Primary Industries –
Fisheries Final Report Series
No. 101
ISSN 1449-9967



NSW DEPARTMENT OF
PRIMARY INDUSTRIES



Recreational
Fishing Trusts

Reproduction and growth of dusky flathead (*Platycephalus fuscus*) in NSW estuaries

July 2008

Authors: Gray, C.A. & Barnes, L.M.
Published By: NSW Department of Primary Industries (now incorporating NSW Fisheries)
Postal Address: Cronulla Fisheries Research Centre of Excellence, PO Box 21, NSW, 2230, Australia
Internet: www.dpi.nsw.gov.au

© NSW Department of Primary Industries and the Fisheries

This work is copyright. Except as permitted under the Copyright Act, no part of this reproduction may be reproduced by any process, electronic or otherwise, without the specific written permission of the copyright owners. Neither may information be stored electronically in any form whatsoever without such permission.

DISCLAIMER

The publishers do not warrant that the information in this report is free from errors or omissions. The publishers do not accept any form of liability, be it contractual, tortuous or otherwise, for the contents of this report for any consequences arising from its use or any reliance placed on it. The information, opinions and advice contained in this report may not relate to, or be relevant to, a reader's particular circumstance.

ISSN 1449-9967

(Note: Prior to July 2004, this report series was published as the 'NSW Fisheries Final Report Series' with ISSN number 1440-3544)

TABLE OF CONTENTS

Table of Contents	i
List of Tables.....	ii
List of Figures	ii
Acknowledgements.....	iii
Non-Technical Aummary	iv
1. INTRODUCTION	5
2. OBJECTIVES	5
3. METHODS.....	6
3.1. <i>Sampling</i>	6
3.2. <i>Reproductive biology</i>	6
3.3. <i>Age and growth</i>	9
4. RESULTS	11
4.1. <i>Reproduction</i>	11
4.2. <i>Age and growth</i>	15
5. DISCUSSION	17
5.1. <i>Reproduction</i>	17
5.2. <i>Age and growth</i>	18
5.3. <i>Management implications</i>	18
6. LITERATURE CITED.....	19

LIST OF TABLES

Table 1.1.	The stages of gonad (ovary and teste) development used to macroscopically classify the sexual maturity of dusky flathead.....	8
Table 1.2.	Summary calculations for male and female L_{50} estimates between November and March.....	12
Table 1.3.	Summary calculations for male and female A_{50} estimates between November and March.	12

LIST OF FIGURES

Figure 1.1.	Location of estuaries in NSW where dusky flathead were sampled.	7
Figure 1.2.	Photo showing different staged female gonads	8
Figure 1.3.	Photo of a sectioned otolith showing alternate zones of opaque and translucent material.....	10
Figure 1.4.	Mean monthly gonadosomatic indices (± 1 S.E) for male and female dusky flathead equal to or larger than the estimated L_{50}	12
Figure 1.5.	Estimated length at maturity (L_{50}) of dusky flathead	13
Figure 1.6.	Estimated age at maturity (A_{50}) of dusky flathead	14
Figure 1.7.	Length-weight relationships for male and female dusky flathead.....	15
Figure 1.8.	Estimated growth of dusky flathead.	16

ACKNOWLEDGEMENTS

This study was initially funded by the NSW Department of Primary Industries. The Recreational Fishing Saltwater Trust Fund provided additional funding for the completion of the project. The latter funded the sampling between 2004 and 2006 and the analyses of data and compilation of the report.

Numerous staff at the Cronulla Fisheries Research Centre provided assistance with sampling and processing of samples and the capture and maintenance of fish held within the Aquarium facility; notably Mr Damian Young, Mr Darren Reynolds, Mr Dylan Van der Meulen and Mr Ben Brewer.

Dr Kevin Rowling and Mr Ben Kendall provided constructive comments on the report.

We thank the various fishers who assisted with sampling and provided samples of flathead.

NON-TECHNICAL SUMMARY

Reproduction and growth of dusky flathead (*Platycephalus fuscus*) in NSW estuaries

OBJECTIVE:

Determine the reproductive cycle, length and age at maturity and the timing and location of spawning of dusky flathead in NSW.

SUMMARY:

Dusky flathead is an important finfish species harvested by recreational and commercial fishers in NSW. Because of concerns over the long-term sustainability of the resource in NSW, the minimum legal length (MLL) of dusky flathead was increased from 33 to 36 cm total length (TL) in July 2001, with further increases (to 40 cm TL) proposed. The initial change in minimum legal length was based on available information concerning the length that dusky flathead spawn. However, detailed information of the reproductive biology of dusky flathead from NSW was lacking. This study was the first step to redress this situation.

Dusky flathead were sampled from various locations throughout NSW between 2001 and 2006 to investigate aspects of their biology. The study identified that female dusky flathead have an extended reproductively active period between November and March while male reproductive activity occurs mainly between September and March. Female dusky flathead in spawning condition were caught only near the entrance of estuaries and in adjacent coastal waters. It was determined that the length at which 50 percent of the population is mature (L_{50}) during periods of peak reproductive activity was 31.72 cm TL for males and 56.75 cm TL for females. The corresponding estimated age at which 50 percent of the population is reproductively mature (A_{50}) during these same periods was 1.22 years for males and 4.55 years for females.

Females grew faster and attained a greater overall maximum TL, weight and age than males. The largest observed female was 98.5cm TL (7.5 kg), and the oldest was estimated to be 16 years, whereas the largest male was 61.5cm TL (weight) and 11 years of age. Although the growth rates and maximum attained lengths varied between sexes, the relationship between length and weight of fish did not differ between sexes.

The current MLL of 36cm TL appears to protect approximately 5 % of the female and 75% of the male spawning population. To provide greater protection to mature females so to allow 50% of females to spawn at least once would require an increase in the MLL to 57cm TL. Such an increase would virtually eliminate males from the fishery. Many other factors, such as the survival rates of released fish from recreational and commercial fishing gears, need to be considered in critiquing future management options, including changes in the MLL, for this species.

1. INTRODUCTION

The dusky flathead (*Platycephalus fuscus*) is endemic to Australia, inhabiting estuaries and nearshore coastal waters along the east coast between Cairns in Queensland and the Gippsland Lakes in Victoria (Kailola *et al.* 1993). Few studies have investigated the life history characteristics of dusky flathead, even though it is a key component of fisheries throughout its distribution, including the estuarine recreational and commercial fisheries in New South Wales (NSW) (Kerby and Brown 1994, Gray *et al.* 2002, 2005, Steffe *et al.* 2005a, b).

Dusky flathead are bottom dwelling ambush predators that normally occur on soft substrates, including mud, sand and seagrass. Tag-release studies show that dusky flathead are capable of moving between estuaries and that they can grow relatively fast (West 1993 and NSW DPI unpublished data). They can potentially attain 18 cm total length (TL) in 1 year, and 40 cm TL in 3 years (Dredge 1976, West 1993, Gray *et al.* 2002); with their maximum reported length being 120 cm TL (approximately 15 kg) (Kailola *et al.* 1993). Reported spawning times of dusky flathead vary between September to March in northern Queensland, November to February in southern Queensland, and January to March in NSW and Victoria (Dredge 1976, SPCC 1981, Russell 1988). There is also conflicting and variable information concerning length at maturity, which has been reported to vary from 46 cm (male) and 56 cm (female) in Queensland to 26 cm (male and female) in southern NSW and Victoria (see Kailola *et al.* 1993). Debate also surrounds whether dusky flathead are protandrous sex reversers (i.e., change sex from male to female) (Dredge 1976, SPCC 1981).

Concerns over the status of populations of dusky flathead in NSW (see Gray *et al.* 2002) led to the minimum legal length (MLL) of dusky flathead increasing from 33 to 36 cm TL in July 2001. Further increases and restrictions in minimum and maximum length at capture have been proposed. At present however, there is not enough information on the reproductive biology of the species in NSW to informatively debate the proposed options. Greater information concerning the reproductive biology of dusky flathead, along with knowledge of their age and growth characteristics, is vital to developing more appropriate management strategies for this species.

This study aimed to redress the current lack of biological information on dusky flathead in NSW by investigating aspects of their reproductive biology, age and growth dynamics.

2. OBJECTIVES

Determine: (1) length and age at sexual maturity; (2) timing and location of spawning; and (3) age and growth characteristics, of dusky flathead in NSW.

3. METHODS

3.1. Sampling

Dusky flathead were sampled in 2001 and 2002 from commercial catches in the Clarence River, Wallis Lake, Tuggerah Lake and Lake Illawarra. Because the majority of samples were restricted to legal sized fish, there was a need to sample a greater length range of fishes, particularly undersized individuals. Further sampling was conducted between 2004 and 2006 in the Clarence River, Lake Macquarie, Tuggerah Lakes, St Georges Basin and the Tuross River (Figure 1.1) to collect undersized and legal sized individuals. A variety of research sampling gears including gillnets, seine nets and angling techniques were employed to sample dusky flathead populations. Additional samples were obtained from the non-retained portion of commercial catches. Samples were also collected from coastal waters adjacent to the Clarence River by chartered commercial prawn trawler operators between November 2005 and March 2007.

The date and location of capture, total length (0.1 cm) and total weight (0.1 g), of each retained dusky flathead was recorded. Fish were then dissected for reproductive investigation and sagittal otoliths were removed, cleaned, dried and stored in labelled paper envelopes before being processed for age determination.

3.2. Reproductive biology

Each retained fish was dissected and its sex determined by macroscopic examination of the gonads (ovaries and testes). When gonads lacked development (i.e., not able to distinguish sex), the fish was classified as juvenile. Macroscopically examined gonads were assigned a reproductive developmental stage based on size, colour and texture, and for females the visibility of oocytes, according to the criteria described in Scott and Pankhurst (1992) (see Table 1.1 and Figure 1.2). The gonads of each fish were then removed and weighed (0.1 g) so that the Gonadosomatic Index (GSI) could be calculated:

$$GSI = (\text{gonad weight} / \text{whole fish weight}) * 100$$

Changes in GSI values throughout time were used to estimate the spawning season; elevated GSI values were interpreted as indicating reproductive activity.

During periods of observed reproductive activity, fish with gonads staged III to VII were considered mature (i.e., capable of spawning during the current reproductive season), whereas individuals staged I and II were considered immature (i.e., not capable of spawning during the current reproductive season). The proportion of fish that were mature within each 1 cm length category and each yearly age class was calculated, and a general least squares modelling approach was used to fit a logistic curve to the data using R statistical programming language. Female maturity calculations were determined from samples pooled across all locations and sampling periods to determine the length (L_{50}) and age (A_{50}) at which 50 % of the sampled population were mature. Maturity calculations for male flathead were pooled across locations and between 2004 and 2006 to determine the length (L_{50}) and age (A_{50}) at which 50 % of the sampled population were mature. The temporal scale analysed was restricted to males sampled between 2004 and 2006 due to potential misclassification of male gonads in the earlier sampling periods.

4. RESULTS

A total of 77,83 dusky flathead, comprising 394 juveniles, 1,745 males and 5,644 females, ranging from 5.7 to 98.5 cm TL were examined throughout the study.

4.1. Reproduction

Female dusky flathead were reproductively active primarily for 5 months between November and March while males were reproductively active for 7 months between September and March as indicated by an increase in the observed GSI (Figure 1.4). The maximum recorded GSI for males and females was 4.7% and 16.6% respectively. No females and only 3 males were observed in reproductive condition for the 3 months between June and August. Females with hydrated oocytes and ovulating individuals were collected from sites close to the entrance of the Clarence River, Wallis Lake, Tuggerah, Lake Illawarra and Tuross River.

The length at which 50% ($L_{50} \pm 1$ SE) of individuals sampled in estuaries attained sexual maturity during periods of peak reproductive activity was estimated to be 56.75 ± 0.6 cm TL for female and 31.72 ± 1.08 cm TL for male dusky flathead (Figure 1.5, Table 1.2). The estimated age at which 50% ($A_{50} \pm 1$ SE) of individuals attained sexual maturity for females was 4.55 ± 0.13 years and 1.22 ± 0.44 years for males (Figure 1.6, Table 1.3). The smallest observed mature female and male was 22.3 cm and 18.8 cm TL respectively. Less than 5 % of females, but approximately 75 % of males, were classified mature at the current MLL of 36cm TL during the spawning period.

Ovarian and testicular tissue was never observed together in the same gonad throughout extensive macroscopic examination, even though gonad samples were examined from fish with varying lengths, ages, multiple locations and over several years of sampling.




Figure 1.4. Mean monthly gonadosomatic indices (± 1 S.E) for male and female dusky flathead equal to or larger than the estimated L_{50} .

Table 1.2. Summary calculations for male and female L_{50} estimates between November and March.

	Male	Female
L_{50} (cm)	31.72	56.75
Standard Error (cm)	1.08	0.6

Table 1.3. Summary calculations for male and female A_{50} estimates between November and March.

	Male	Female
A_{50} (Years)	1.22	4.55
Standard Error (Years)	0.44	0.13




Figure 1.5. Estimated length at maturity (L_{50}) of dusky flathead: a) Male 31.72 ± 1.08 cm ($n = 484$), b) Female 56.75 ± 0.6 cm ($n = 3,371$). The current minimum legal limit (MLL) is 36 cm TL.




Figure 1.6. Estimated age at maturity (A_{50}) of dusky flathead: a) Male 1.22 ± 0.44 years ($n=190$), b) Female 4.55 ± 0.13 years ($n = 2,288$).

4.2. Age and growth

The length-weight relationship between male and female dusky flathead was very similar (Figure 1.7). The length-weight equation for males and females was:

$$\text{Males: Weight} = 2.86 \times 10^{-3} * \text{Total Length}^{(3.213)};$$

$$\text{Females: Weight} = 2.09 \times 10^{-3} * \text{Total Length}^{(3.283)}$$




Figure 1.7. Length-weight relationships for male and female dusky flathead.

Female dusky flathead grew faster and attained a greater maximum length and age compared to males (Figure 1.8). The largest female sampled was 98.5 cm TL, weighing 7.5 kilograms and an estimated age of 13 years. The oldest female was estimated to be 16 years (88.5cm TL). The largest male sampled was 61.5 cm TL, weighing 1.58 kg and an estimated age of 11 years.

There was considerable variation in length at age for both sexes (Figure 1.8). The parameter estimates of the von Bertalanffy growth curves differed between sexes. For fish greater than 2 years old, the estimated mean length at age was consistently greater for females than for males.





Figure 1.8. Estimated growth of a) male (n = 1,421) and b) female (n = 4,333) dusky flathead.

5. DISCUSSION

5.1. Reproduction

The predominant spawning period we observed for female dusky flathead was between November and March while the observed male spawning period extended from September to March. This general timing of spawning concurs with previous reports from Queensland (September to March in northern Queensland and November to February in southern Queensland). It is however, more extended than that previously reported for NSW (January to March) (Dredge 1976, SPCC 1981, Russell 1988). Larvae of dusky flathead have been captured in estuaries and coastal waters of NSW between September and May (Miskiewicz 1987, Gray & Miskiewicz 2000).

Estimates of fecundity for dusky flathead range from 294,000 to 3,948,000 eggs (Brown *et al.* 1994) however, the mode of spawning is unclear. *P. endrachtensis* and *P. indicus* exhibit an extended spawning period and exhibit bimodal oocyte diameter frequency distributions which lead Lewis (1971) to suggest these species have multiple spawning events throughout the year. Similarly Hyndes *et al.* (1992b) found *P. speculator* to be multiple spawners between December and March. Additional investigation is required to determine the spawning mode of dusky flathead.

Dusky flathead were observed to be in spawning condition in the marine dominated lower reaches of estuaries, ovulating females were sampled from the Clarence River, Wallis Lake, Tuggerah Lake, Lake Illawarra and Tuross River at sites close to the mouth of the estuary. These observations support the general hypothesis that dusky flathead spawn in the lower reaches of estuaries and in nearshore coastal waters (SPCC 1981, Kailola *et al.* 1993). Furthermore, in St Georges Basin and Lake Macquarie where samples were collected upstream of the mouth, only females with vitellogenic oocytes (i.e., not in spawning condition) were observed.

Our estimated L_{50} of 56.75 cm TL for females was similar to the estimate for Queensland of 56cm TL (Russell 1998) but larger than the previously estimated 38cm TL in Botany Bay, New South Wales (SPCC 1981). In contrast however, our estimated 31.72cm TL length at maturity for males was considerably less than that estimated for Queensland (46cm TL), but was similar to that reported by Winstanley (1985) and SPCC (1981) of 26 and 32 cm TL respectively. The estimated L_{50} for male dusky flathead appears to be highly variable. In other teleost species, it has been found that social and behavioural cues, as well as differential hormonal expression influence the timing of male maturation (Oliveira *et al.* 2005, Kuwamura *et al.* 2007). Further investigation is required to determine the roles of these factors in male dusky flathead maturation.

Female dusky flathead reach sexual maturity later and at a larger size than males. On average, males take 1.2 years to reach the L_{50} , females however, take more than 4 years to reach the estimated L_{50} although some females mature at approximately 2 years of age. This pattern is typical of many platycephalid species (Lewis 1971, Jordon 2001) and may be related to an increase in female reproductive potential at larger sizes as seen for many teleost species (Hughes and Stewart 2006, McDermott *et al.* 2007).

Several platycephalids are known to exhibit protandrous hermaphroditism (Fujii 1971) and it has been argued that dusky flathead undergo protandrous sex change, where individuals first function as males for several years before changing sex (Dredge 1976). This has been based on the observed skewed sex ratios in dusky flathead populations with the smaller size classes being dominated by males and the larger size classes by females. No hermaphroditic fish were observed in this study or in previous studies in NSW (Gray *et al.* 2002, West 1993), suggesting that dusky flathead do not

change sex, but rather male fish do not grow as large or as fast as females. *P. indicus* (Masuda *et al.* 2000) and *P. bassensis* (Jordan 1998) show the same dimorphic growth pattern between sexes with no evidence of protandrous sex change.

5.2. Age and growth

Dusky flathead grow relatively quickly, with fish estimated to reach a mean length of approximately 30 cm TL in 1 year and 42 and 59 cm TL after 5 years for males and females respectively. It is evident that growth rate is highly variable between individuals. Our estimated rates of growth are consistent with previous growth estimates (Dredge 1976, West 1993, Gray *et al.* 2002). We estimate that most dusky flathead are approximately 2 years old when they reach the current MLL of 36 cm TL.

As with other platycephalids, females grew faster and attained a larger maximum length than males. However, the maximum length (and weight) is considerably greater for dusky flathead than for other Platycephalids such as *P. bassensis* (Jordan 1998), *P. caeruleopunctatus*, *P. speculator* (Brown 1977) and *P. richardsoni* (Cui *et al.* 2005). Our estimate of longevity of dusky flathead of 16 years is comparable to other species of flathead, such as *P. richardsoni* and *P. bassensis*, although these species attain a smaller maximum length than dusky flathead.

The female von Bertalanffy growth model maximum size of 127cm was similar to the largest reported dusky flathead reported (120cm) by Kailola (1993). The largest female dusky flathead sampled within the current study was 98.5cm considerably less than the maximum observed or modelled length. The modelled maximum male length of 44cm was smaller than the 61.5cm largest male observed. The poorer fit of the von Bertalanffy growth model to the male growth may be due to the rarity of larger and older male fishes within the sample. Similarly, larger males may have not been sampled as effectively as smaller individuals due to factors such as habitat association or behavioural characteristics. In general however, larger fishes of both sexes do not appear to be as abundant as smaller individuals nor were they as large as has previously been observed (Kailola 1993) within the habitats and locations that were sampled during this study.

5.3. Management implications

The current MLL of 36 cm TL provides significant protection for mature male dusky flathead, but not for mature females. Approximately 5% of females are estimated to be mature at 36 cm TL. The MLL for many species is set at the female L_{50} ; for dusky flathead this would be 57 cm TL. Such an increase in the MLL of dusky flathead would virtually eliminate males from exploitation. At present, there is an additional management regulation of only 1 dusky flathead > 70cm TL to be retained by recreational anglers per day to provide greater protection to large females.

In assessing the implications of the current research on any future changes to the MLL or management intervention, an overall management objective for the population must be formulated. Examples of objectives on which to base management decisions upon may include, but are not limited to, the opportunity to spawn once before recruiting into the fishery, ensuring sufficient reproduction to maintain stocks, maximising value of the resource and maximising yield (Stewart and Hughes 2008). An assessment of dusky flathead stocks, as well as other key species of flathead, needs to be made in order to provide appropriate information on which management objectives can be based and ultimately implemented.

6. LITERATURE CITED

- Broadhurst, M.K., Gray, C.A., Young, D.J., Johnson, D.D. (2003) Relative efficiency and size selectivity of bottom-set gillnets for dusky flathead, *Platycephalus fuscus* and other species in New South Wales, Australia. *Archive of Fishery and Marine Research* 50, 289–302.
- Brown, I.W. (1977) Ecology of Three Sympatric Flatheads (Platycephalidae) in Port Phillip Bay, Victoria. PhD Thesis Monash University, Victoria.
- Cui, G., Knuckey, I. and Klaer, N. (2005) Tiger flathead stock assessment using data up to 2004. Report to SESSF RAG Workshop, August 2005. CSIRO Marine Research, Hobart.
- Dredge, M.C.L. (1976) Aspects of the ecology of three estuarine dwelling fish in south-east Queensland. M.Sc. Thesis, University of Queensland. 122 pp.
- Gray, C.A. (2002) Management implications of discarding in an estuarine multi-species gill net fishery. *Fisheries Research* 56, 177–192.
- Gray, C.A. (2004) An overview of commercial fishing in Lake Illawarra: trends in reported catch and effort, bycatch and discarding. *Wetlands (Australia)* 21, 152–162.
- Gray, C.A., Gale, V.J., Stringfellow, S.L., Raines, L.P. (2002) Variations in sex, length and age compositions of commercial catches of *Platycephalus fuscus* (Pisces: Platycephalidae) in New South Wales, Australia. *Marine and Freshwater Research* 53, 1091–1100.
- Gray, C.A., Johnson, D.D., Young, D.J., Broadhurst, M.K. (2004) Discards from the commercial gillnet fishery for dusky flathead, *Platycephalus fuscus*, in New South Wales, Australia: spatial variability and initial effects of change in minimum legal length of target species. *Fisheries Management and Ecology* 11, 323–333.
- Gray, C.A., Johnson, D.D., Broadhurst, M.K., Young, D.J. (2005) Seasonal, spatial and gear-related influences on relationships between retained and discarded catches in a multi-species gillnet fishery. *Fisheries Research* 75, 56–72.
- Gray, C.A., Broadhurst, M.K., Johnson, D.D., Young, D.J. (2005) Influences of hanging ratio, fishing height, twine diameter and material of bottom-set gillnets on catches of dusky flathead *Platycephalus fuscus* and non-target species in New South Wales, Australia. *Fisheries Science* 71, 1216–1227.
- Gray, C.A., Miskiewicz, A.G. (2000) Larval fish assemblages in south-east Australian coastal waters: seasonal and spatial structure. *Estuarine, Coastal and Shelf Science* 50, 549–570.
- Hughes, J.A. and Stewart, J. (2006) Reproductive biology of three commercially important Hemiramphid species in south-eastern Australia. *Environmental Biology of Fishes* 75:237–256.
- Hyndes, G.A., Loneragan, N.R., Potter, I.C. (1992a) Influence of sectioning otoliths on marginal increment trends and age and growth estimates for the flathead *Platycephalus speculator*. *Fishery Bulletin* 90, 276–284.
- Hyndes, G.A., Neirra, F.J., Potter, I.C. (1992b) Reproductive biology and early life history of the marine teleost *Platycephalus speculator* Klunzinger (Platycephalidae) in a temperate Australian estuary. *Journal of Fish Biology* 40, 859–874.

- Jordan, A.R. (2001) Reproductive biology, early life history and settlement distribution of sand flathead (*Platycephalus bassensis*) in Tasmania. *Marine and Freshwater Research* 52, 589–601.
- Kailola, P.J., Williams, M.J., Stewart, P.C., Reichelt, R.E., McNee, A., Grieve, C. (1993) Australian Fisheries Resources. Bureau of Resource Sciences, Department of Primary Industries and Energy, and Fisheries Research and Development Corporation. Canberra, Australia. 422 pp.
- Kerby, B.M., Brown, I.W. (1994) Bream, whiting and flathead in south-east Queensland: a review of the literature. Queensland Department of Primary Industries, QI94028.
- Lewis, A.D. (1971) A comparative study of the biology of five south Queensland flatheads (Platycephalidae). Honours Thesis University of Queensland.
- McDermott, S.F., Maslenikov, K.P., Gunderson, D.R. (2007) Annual fecundity, batch fecundity, and oocyte atresia of Atka mackerel (*Pleurogrammus monopterygius*) in Alaskan waters. *Fisheries Bulletin* 105:19–29.
- Miskiewicz, A.G. (1987) Taxonomy and ecology of fish larvae in Lake Macquarie and New South Wales coastal waters. PhD thesis, University of New South Wales.
- Russell, D.J. (1998) An assessment of the east Queensland inshore gillnet fishery. *Queensland Department of Primary Industries, Information Series Q188024*.
- SPCC. (1981) Ecology of Fish in Botany Bay – Biology of commercially and recreationally valuable species. State Pollution Control Commission BBS 23B.
- Steffe, A.S., Murphy, J.J., Chapman, D.J., Barrett, G.P., Gray, C.A. (2005) An assessment of changes in the daytime, boat-based, recreational fishery of the Tuross Lake estuary following the establishment of a ‘Recreational Fishing Haven’. NSW Department of Primary Industries – Fisheries Final Report Series No. 81. ISSN 1449-9967.
- Steffe, A.S., Murphy, J.J., Chapman, D.J., Gray, C.A. (2005) An assessment of changes in the daytime, boat-based, recreational fishery of Lake Macquarie following the establishment of a ‘Recreational Fishing Haven’. NSW Department of Primary Industries – Fisheries Final Report Series No. 79. ISSN 1449-9967.
- Stewart, J and Hughes, J.M. (2008) Determining appropriate sizes at harvest for species shared by the commercial trap and recreational fisheries in New South Wales. NSW Department of Primary Industries – Fisheries Final Report Series No. 97. ISSN 1449-9967.
- von Bertalanffy, L. (1938) A quantitative theory of organic growth. *Human Biology* 10:181–213
- West, R.J. (1993) The fisheries resources of two southeastern Australian rivers. Unpublished PhD Thesis, University of New South Wales, Australia.

Other titles in this series:**ISSN 1440-3544 (NSW Fisheries Final Report Series)**

- No. 1 Andrew, N.L., Graham, K.J., Hodgson, K.E. and Gordon, G.N.G., 1998. Changes after 20 years in relative abundance and size composition of commercial fishes caught during fishery independent surveys on SEF trawl grounds. Final Report to Fisheries Research and Development Corporation. Project No. 96/139.
- No. 2 Virgona, J.L., Deguara, K.L., Sullings, D.J., Halliday, I. and Kelly, K., 1998. Assessment of the stocks of sea mullet in New South Wales and Queensland waters. Final Report to Fisheries Research and Development Corporation. Project No. 94/024.
- No. 3 Stewart, J., Ferrell, D.J. and Andrew, N.L., 1998. Ageing Yellowtail (*Trachurus novaezelandiae*) and Blue Mackerel (*Scomber australasicus*) in New South Wales. Final Report to Fisheries Research and Development Corporation. Project No. 95/151.
- No. 4 Pethebridge, R., Lugg, A. and Harris, J., 1998. Obstructions to fish passage in New South Wales South Coast streams. Final report to Cooperative Research Centre for Freshwater Ecology. 70pp.
- No. 5 Kennelly, S.J. and Broadhurst, M.K., 1998. Development of by-catch reducing prawn-trawls and fishing practices in NSW's prawn-trawl fisheries (and incorporating an assessment of the effect of increasing mesh size in fish trawl gear). Final Report to Fisheries Research and Development Corporation. Project No. 93/180. 18pp + appendices.
- No. 6 Allan, G.L. and Rowland, S.J., 1998. Fish meal replacement in aquaculture feeds for silver perch. Final Report to Fisheries Research and Development Corporation. Project No. 93/120-03. 237pp + appendices.
- No. 7 Allan, G.L., 1998. Fish meal replacement in aquaculture feeds: subprogram administration. Final Report to Fisheries Research and Development Corporation. Project No. 93/120. 54pp + appendices.
- No. 8 Heasman, M.P., O'Connor, W.A. and O'Connor, S.J., 1998. Enhancement and farming of scallops in NSW using hatchery produced seedstock. Final Report to Fisheries Research and Development Corporation. Project No. 94/083. 146pp.
- No. 9 Nell, J.A., McMahon, G.A. and Hand, R.E., 1998. Tetraploidy induction in Sydney rock oysters. Final Report to Cooperative Research Centre for Aquaculture. Project No. D.4.2. 25pp.
- No. 10 Nell, J.A. and Maguire, G.B., 1998. Commercialisation of triploid Sydney rock and Pacific oysters. Part 1: Sydney rock oysters. Final Report to Fisheries Research and Development Corporation. Project No. 93/151. 122pp.
- No. 11 Watford, F.A. and Williams, R.J., 1998. Inventory of estuarine vegetation in Botany Bay, with special reference to changes in the distribution of seagrass. Final Report to Fishcare Australia. Project No. 97/003741. 51pp.
- No. 12 Andrew, N.L., Worthington D.G., Brett, P.A. and Bentley N., 1998. Interactions between the abalone fishery and sea urchins in New South Wales. Final Report to Fisheries Research and Development Corporation. Project No. 93/102.
- No. 13 Jackson, K.L. and Ogburn, D.M., 1999. Review of depuration and its role in shellfish quality assurance. Final Report to Fisheries Research and Development Corporation. Project No. 96/355. 77pp.
- No. 14 Fielder, D.S., Bardsley, W.J. and Allan, G.L., 1999. Enhancement of Mulloway (*Argyrosomus japonicus*) in intermittently opening lagoons. Final Report to Fisheries Research and Development Corporation. Project No. 95/148. 50pp + appendices.
- No. 15 Otway, N.M. and Macbeth, W.G., 1999. The physical effects of hauling on seagrass beds. Final Report to Fisheries Research and Development Corporation. Project No. 95/149 and 96/286. 86pp.
- No. 16 Gibbs, P., McVea, T. and Loudon, B., 1999. Utilisation of restored wetlands by fish and invertebrates. Final Report to Fisheries Research and Development Corporation. Project No. 95/150. 142pp.
- No. 17 Ogburn, D. and Ruello, N., 1999. Waterproof labelling and identification systems suitable for shellfish and other seafood and aquaculture products. Whose oyster is that? Final Report to Fisheries Research and Development Corporation. Project No. 95/360. 50pp.

- No. 18 Gray, C.A., Pease, B.C., Stringfellow, S.L., Raines, L.P. and Walford, T.R., 2000. Sampling estuarine fish species for stock assessment. Includes appendices by D.J. Ferrell, B.C. Pease, T.R. Walford, G.N.G. Gordon, C.A. Gray and G.W. Liggins. Final Report to Fisheries Research and Development Corporation. Project No. 94/042. 194pp.
- No. 19 Otway, N.M. and Parker, P.C., 2000. The biology, ecology, distribution, abundance and identification of marine protected areas for the conservation of threatened Grey Nurse Sharks in south east Australian waters. Final Report to Environment Australia. 101pp.
- No. 20 Allan, G.L. and Rowland, S.J., 2000. Consumer sensory evaluation of silver perch cultured in ponds on meat meal based diets. Final Report to Meat & Livestock Australia. Project No. PRCOP.009. 21pp + appendices.
- No. 21 Kennelly, S.J. and Scandol, J. P., 2000. Relative abundances of spanner crabs and the development of a population model for managing the NSW spanner crab fishery. Final Report to Fisheries Research and Development Corporation. Project No. 96/135. 43pp + appendices.
- No. 22 Williams, R.J., Watford, F.A. and Balashov, V., 2000. Kooragang Wetland Rehabilitation Project: History of changes to estuarine wetlands of the lower Hunter River. Final Report to Kooragang Wetland Rehabilitation Project Steering Committee. 82pp.
- No. 23 Survey Development Working Group, 2000. Development of the National Recreational and Indigenous Fishing Survey. Final Report to Fisheries Research and Development Corporation. Project No. 98/169. (Volume 1 – 36pp + Volume 2 – attachments).
- No.24 Rowling, K.R and Raines, L.P., 2000. Description of the biology and an assessment of the fishery of Silver Trevally *Pseudocaranx dentex* off New South Wales. Final Report to Fisheries Research and Development Corporation. Project No. 97/125. 69pp.
- No. 25 Allan, G.L., Jantrarotai, W., Rowland, S., Kosuturak, P. and Booth, M., 2000. Replacing fishmeal in aquaculture diets. Final Report to the Australian Centre for International Agricultural Research. Project No. 9207. 13pp.
- No. 26 Gehrke, P.C., Gilligan, D.M. and Barwick, M., 2001. Fish communities and migration in the Shoalhaven River – Before construction of a fishway. Final Report to Sydney Catchment Authority. 126pp.
- No. 27 Rowling, K.R. and Makin, D.L., 2001. Monitoring of the fishery for Gemfish *Rexea solandri*, 1996 to 2000. Final Report to the Australian Fisheries Management Authority. 44pp.
- No. 28 Otway, N.M., 1999. Identification of candidate sites for declaration of aquatic reserves for the conservation of rocky intertidal communities in the Hawkesbury Shelf and Batemans Shelf Bioregions. Final Report to Environment Australia for the Marine Protected Areas Program. Project No. OR22. 88pp.
- No. 29 Heasman, M.P., Goard, L., Diemar, J. and Callinan, R., 2000. Improved Early Survival of Molluscs: Sydney Rock Oyster (*Saccostrea glomerata*). Final report to the Aquaculture Cooperative Research Centre. Project No. A.2.1. 63pp.
- No. 30 Allan, G.L., Dignam, A and Fielder, S., 2001. Developing Commercial Inland Saline Aquaculture in Australia: Part 1. R&D Plan. Final Report to Fisheries Research and Development Corporation. Project No. 1998/335.
- No. 31 Allan, G.L., Banens, B. and Fielder, S., 2001. Developing Commercial Inland Saline Aquaculture in Australia: Part 2. Resource Inventory and Assessment. Final report to Fisheries Research and Development Corporation. Project No. 1998/335. 33pp.
- No. 32 Bruce, A., Grows, I. and Gehrke, P., 2001. Woronora River Macquarie Perch Survey. Final report to Sydney Catchment Authority, April 2001. 116pp.
- No. 33 Morris, S.A., Pollard, D.A., Gehrke, P.C. and Pogonoski, J.J., 2001. Threatened and Potentially Threatened Freshwater Fishes of Coastal New South Wales and the Murray-Darling Basin. Report to Fisheries Action Program and World Wide Fund for Nature. Project No. AA 0959.98. 177pp.
- No. 34 Heasman, M.P., Sushames, T.M., Diemar, J.A., O'Connor, W.A. and Foulkes, L.A., 2001. Production of Micro-algal Concentrates for Aquaculture Part 2: Development and Evaluation of Harvesting, Preservation, Storage and Feeding Technology. Final Report to Fisheries Research and Development Corporation. Project No. 1993/123 and 1996/342. 150pp + appendices.
- No. 35 Stewart, J. and Ferrell, D.J., 2001. Mesh selectivity in the NSW demersal trap fishery. Final Report to Fisheries Research and Development Corporation. Project No. 1998/138. 86pp.

- No. 36 Stewart, J., Ferrell, D.J., van der Walt, B., Johnson, D. and Lowry, M., 2001. Assessment of length and age composition of commercial kingfish landings. Final Report to Fisheries Research and Development Corporation. Project No. 1997/126. 49pp.
- No. 37 Gray, C.A. and Kennelly, S.J., 2001. Development of discard-reducing gears and practices in the estuarine prawn and fish haul fisheries of NSW. Final Report to Fisheries Research and Development Corporation. Project No. 1997/207. 151pp.
- No. 38 Murphy, J.J., Lowry, M.B., Henry, G.W. and Chapman, D., 2002. The Gamefish Tournament Monitoring Program – 1993 to 2000. Final report to Australian Fisheries Management Authority. 93pp.
- No. 39 Kennelly, S.J. and McVea, T.A. (Ed), 2002. Scientific reports on the recovery of the Richmond and Macleay Rivers following fish kills in February and March 2001. 325pp.
- No. 40 Pollard, D.A. and Pethebridge, R.L., 2002. Report on Port of Botany Bay Introduced Marine Pest Species Survey. Final Report to Sydney Ports Corporation. 69pp.
- No. 41 Pollard, D.A. and Pethebridge, R.L., 2002. Report on Port Kembla Introduced Marine Pest Species Survey. Final Report to Port Kembla Port Corporation. 72pp.
- No. 42 O'Connor, W.A, Lawler, N.F. and Heasman, M.P., 2003. Trial farming the akoya pearl oyster, *Pinctada imbricata*, in Port Stephens, NSW. Final Report to Australian Radiata Pty. Ltd. 170pp.
- No. 43 Fielder, D.S. and Allan, G.L., 2003. Improving fingerling production and evaluating inland saline water culture of snapper, *Pagrus auratus*. Final Report to the Aquaculture Cooperative Research Centre. Project No. C4.2. 62pp.
- No. 44 Astles, K.L., Winstanley, R.K., Harris, J.H. and Gehrke, P.C., 2003. Experimental study of the effects of cold water pollution on native fish. A Final Report for the Regulated Rivers and Fisheries Restoration Project. 55pp.
- No. 45 Gilligan, D.M., Harris, J.H. and Mallen-Cooper, M., 2003. Monitoring changes in the Crawford River fish community following replacement of an effective fishway with a vertical-slot fishway design: Results of an eight year monitoring program. Final Report to the Cooperative Research Centre for Freshwater Ecology. 80pp.
- No. 46 Pollard, D.A. and Rankin, B.K., 2003. Port of Eden Introduced Marine Pest Species Survey. Final Report to Coasts & Clean Seas Program. 67pp.
- No. 47 Otway, N.M., Burke, A.L., Morrison, NS. and Parker, P.C., 2003. Monitoring and identification of NSW Critical Habitat Sites for conservation of Grey Nurse Sharks. Final Report to Environment Australia. Project No. 22499. 62pp.
- No. 48 Henry, G.W. and Lyle, J.M. (Ed), 2003. The National Recreational and Indigenous Fishing Survey. Final Report to Fisheries Research and Development Corporation. Project No. 1999/158. 188 pp.
- No. 49 Nell, J.A., 2003. Selective breeding for disease resistance and fast growth in Sydney rock oysters. Final Report to Fisheries Research and Development Corporation. Project No. 1996/357. 44pp. (Also available – a CD-Rom published in March 2004 containing a collection of selected manuscripts published over the last decade in peer-reviewed journals).
- No. 50 Gilligan, D. and Schiller, S., 2003. Downstream transport of larval and juvenile fish. A final report for the Natural Resources Management Strategy. Project No. NRMS R7019. 66pp.
- No. 51 Liggins, G.W., Scandol, J.P. and Kennelly, S.J., 2003. Recruitment of Population Dynamacist. Final Report to Fisheries Research and Development Corporation. Project No. 1993/214.05. 44pp.
- No. 52 Steffe, A.S. and Chapman, J.P., 2003. A survey of daytime recreational fishing during the annual period, March 1999 to February 2000, in Lake Macquarie, New South Wales. NSW Fisheries Final Report. 124pp.
- No. 53 Barker, D. and Otway, N., 2003. Environmental assessment of zinc coated wire mesh sea cages in Botany Bay NSW. Final Report to OneSteel Limited. 36pp.
- No. 54 Grouns, I., Astles, A. and Gehrke, P., 2003. Spatial and temporal variation in composition of riverine fish communities. Final Report to Water Management Fund. Project No. SW1 part 2. 24pp.

- No. 55 Gray, C. A., Johnson, D.D., Young, D.J. and Broadhurst, M. K., 2003. Bycatch assessment of the Estuarine Commercial Gill Net Fishery in NSW. Final Report to Fisheries Research and Development Corporation. Project No. 2000/172. 58pp.
- No. 56 Worthington, D.G. and Blount, C., 2003. Research to develop and manage the sea urchin fisheries of NSW and eastern Victoria. Final Report to Fisheries Research and Development Corporation. Project No. 1999/128. 182pp.
- No. 57 Baumgartner, L.J., 2003. Fish passage through a Deelder lock on the Murrumbidgee River, Australia. NSW Fisheries Final Report. 34pp.
- No. 58 Allan, G.L., Booth, M.A., David A.J. Stone, D.A.J. and Anderson, A.J., 2004. Aquaculture Diet Development Subprogram: Ingredient Evaluation. Final Report to Fisheries Research and Development Corporation. Project No. 1996/391. 171pp.
- No. 59 Smith, D.M., Allan, G.L. and Booth, M.A., 2004. Aquaculture Diet Development Subprogram: Nutrient Requirements of Aquaculture Species. Final Report to Fisheries Research and Development Corporation. Project No. 1996/392. 220pp.
- No. 60 Barlow, C.G., Allan, G.L., Williams, K.C., Rowland, S.J. and Smith, D.M., 2004. Aquaculture Diet Development Subprogram: Diet Validation and Feeding Strategies. Final Report to Fisheries Research and Development Corporation. Project No. 1996/393. 197pp.
- No. 61 Heasman, M.H., 2004. Sydney Rock Oyster Hatchery Workshop 8 – 9 August 2002, Port Stephens, NSW. Final Report to Fisheries Research and Development Corporation. Project No. 2002/206. 115pp.
- No. 62 Heasman, M., Chick, R., Savva, N., Worthington, D., Brand, C., Gibson, P. and Diemar, J., 2004. Enhancement of populations of abalone in NSW using hatchery-produced seed. Final Report to Fisheries Research and Development Corporation. Project No. 1998/219. 269pp.
- No. 63 Otway, N.M. and Burke, A.L., 2004. Mark-recapture population estimate and movements of Grey Nurse Sharks. Final Report to Environment Australia. Project No. 30786/87. 53pp.
- No. 64 Creese, R.G., Davis, A.R. and Glasby, T.M., 2004. Eradicating and preventing the spread of the invasive alga *Caulerpa taxifolia* in NSW. Final Report to the Natural Heritage Trust's Coasts and Clean Seas Introduced Marine Pests Program. Project No. 35593. 110pp.
- No. 65 Baumgartner, L.J., 2004. The effects of Balranald Weir on spatial and temporal distributions of lower Murrumbidgee River fish assemblages. Final Report to the Department of Agriculture, Fisheries & Forestry – Australia (National Heritage Trust MD2001 Fishrehab Program). 30pp.
- No. 66 Heasman, M., Diggles, B.K., Hurwood, D., Mather, P., Pirozzi, I. and Dworjanyan, S., 2004. Paving the way for continued rapid development of the flat (angasi) oyster (*Ostrea angasi*) farming in New South Wales. Final Report to the Department of Transport & Regional Services. Project No. NT002/0195. 40pp.

ISSN 1449-9967 (NSW Department of Primary Industries – Fisheries Final Report Series)

- No. 67 Kroon, F.J., Bruce, A.M., Housefield, G.P. and Creese, R.G., 2004. Coastal floodplain management in eastern Australia: barriers to fish and invertebrate recruitment in acid sulphate soil catchments. Final Report to Fisheries Research and Development Corporation. Project No. 1998/215. 212pp.
- No. 68 Walsh, S., Copeland, C. and Westlake, M., 2004. Major fish kills in the northern rivers of NSW in 2001: Causes, Impacts & Responses. NSW Department of Primary Industries – Fisheries Final Report. 55pp.
- No. 69 Pease, B.C. (Ed), 2004. Description of the biology and an assessment of the fishery for adult longfinned eels in NSW. Final Report to Fisheries Research and Development Corporation. Project No. 1998/127. 168pp.
- No. 70 West, G., Williams, R.J. and Laird, R., 2004. Distribution of estuarine vegetation in the Parramatta River and Sydney Harbour, 2000. Final Report to NSW Maritime and the Australian Maritime Safety Authority. 37pp.
- No. 71 Broadhurst, M.K., Macbeth, W.G. and Wooden, M.E.L., 2005. Reducing the discarding of small prawns in NSW's commercial and recreational prawn fisheries. Final Report to the Fisheries

- Research & Development Corporation. Project No. 2001/031. NSW Department of Primary Industries – Fisheries Final Report Series No. 71. 202pp.
- No. 72. Graham, K.J., Lowry, M.B. and Walford, T.R., 2005. Carp in NSW: Assessment of distribution, fishery and fishing methods. NSW Department of Primary Industries – Fisheries Final Report Series No. 72. 88pp.
- No. 73 Stewart, J., Hughes, J.M., Gray, C.A. and Walsh, C., 2005. Life history, reproductive biology, habitat use and fishery status of eastern sea garfish (*Hyporhamphus australis*) and river garfish (*H. regularis ardelio*) in NSW waters. Final report on the Fisheries Research & Development Corporation Project No. 2001/027. 180pp.
- No. 74 Grouns, I. and Gehrke, P., 2005. Integrated Monitoring of Environmental Flows: Assessment of predictive modelling for river flows and fish. NSW Department of Primary Industries – Fisheries Final Report Series No. 74. 33pp.
- No. 75 Gilligan, D., 2005. Fish communities of the Murrumbidgee catchment: Status and trends. Final report to the Murrumbidgee Catchment Management Authority. Project No. BG4_03. 138pp.
- No. 76 Ferrell, D.J., 2005. Biological information for appropriate management of endemic fish species at Lord Howe Island. NSW Department of Primary Industries – Fisheries Final Report Series No. 76. 18 pp.
- No. 77 Gilligan, D., Gehrke, P. and Schiller, C., 2005. Testing methods and ecological consequences of large-scale removal of common carp. Final report to the Water Management Fund – Programs MFW6 and MUR5. 46pp.
- No. 78 Boys, C.A., Esslemont, G. and Thoms, M.C., 2005. Fish habitat and protection in the Barwon-Darling and Paroo Rivers. Final report to the Department of Agriculture, Fisheries and Forestry – Australia (AFFA). 118pp.
- No. 79 Steffe, A.S., Murphy, J.J., Chapman, D.J. and Gray, C.C., 2005. An assessment of changes in the daytime recreational fishery of Lake Macquarie following the establishment of a 'Recreational Fishing Haven'. NSW Department of Primary Industries – Fisheries Final Report Series No. 79. 103pp.
- No. 80 Gannassin, C. and Gibbs, P., 2005. Broad-Scale Interactions Between Fishing and Mammals, Reptiles and Birds in NSW Marine Waters. Final Report for a project undertaken for the NSW Biodiversity Strategy. NSW Department of Primary Industries – Fisheries Final Report Series No. 80. 171pp.
- No. 81 Steffe, A.S., Murphy, J.J., Chapman, D.J., Barrett, G.P. and Gray, C.A., 2005. An assessment of changes in the daytime, boat-based, recreational fishery of the Tuross Lake estuary following the establishment of a 'Recreational Fishing Haven'. NSW Department of Primary Industries – Fisheries Final Report Series No. 81. 70pp.
- No. 82 Silberschnieder, V. and Gray, C.A., 2005. Arresting the decline of the commercial and recreational fisheries for mulloway (*Argyrosomus japonicus*). Final report on the Fisheries Research & Development Corporation Project No. 2001/027. 71pp.
- No. 83 Gilligan, D., 2005. Fish communities of the Lower Murray-Darling catchment: Status and trends. Final report to the Lower Murray Darling Catchment Management Authority. Project No. MD 005.03. 106pp.
- No. 84 Baumgartner, L.J., Reynoldson, N., Cameron, L. and Stanger, J., 2006. Assessment of a Dual-frequency Identification Sonar (DIDSON) for application in fish migration studies. NSW Department of Primary Industries – Fisheries Final Report Series No. 84. 33pp.
- No. 85 Park, T., 2006. FishCare Volunteer Program Angling Survey: Summary of data collected and recommendations. NSW Department of Primary Industries – Fisheries Final Report Series No. 85. 41pp.
- No. 86 Baumgartner, T., 2006. A preliminary assessment of fish passage through a Denil fishway on the Edward River, Australia. Final report to the Lower Murray Darling Catchment Management Authority, Project No. MD524. 23pp.
- No. 87 Stewart, J., 2007. Observer study in the Estuary General sea garfish haul net fishery in NSW. NSW Department of Primary Industries – Fisheries Final Report Series No. 87. 23pp.

- No. 88 Faragher, R.A., Pogonoski, J.J., Cameron, L., Baumgartner, L. and van der Walt, B., 2007. Assessment of a stocking program: Findings and recommendations for the Snowy Lakes Trout Strategy. NSW Department of Primary Industries – Fisheries Final Report Series No. 88. 46pp.
- No. 89 Gilligan, D., Rolls, R., Merrick, J., Lintermans, M., Duncan, P. and Kohen, J., 2007. Scoping knowledge requirements for Murray crayfish (*Euastacus armatus*). Final report to the Murray Darling Basin Commission for Project No. 05/1066 NSW Department of Primary Industries – Fisheries Final Report Series No. 89. 103pp.
- No. 90 Kelleway, J., Williams, R.J. and Allen, C.B., 2007. An assessment of the saltmarsh of the Parramatta River and Sydney Harbour. Final report to NSW Maritime Authority. NSW DPI – Fisheries Final Report Series No. 90. 100pp.
- No. 91 Williams, R.J. and Thiebaud, I., 2007. An analysis of changes to aquatic habitats and adjacent land-use in the downstream portion of the Hawkesbury Nepean River over the past sixty years. Final report to the Hawkesbury-Nepean Catchment Management Authority. NSW DPI – Fisheries Final Report Series No. 91. 97pp.
- No. 92 Baumgartner, L., Reynoldson, N., Cameron, L. and Stanger, J. The effects of selected irrigation practices on fish of the Murray-Darling Basin. Final report to the Murray Darling Basin Commission for Project No. R5006. NSW Department of Primary Industries – Fisheries Final Report Series No. 92. 90pp.
- No. 93 Rowland, S.J., Landos, M., Callinan, R.B., Allan, G.L., Read, P., Mifsud, C., Nixon, M., Boyd, P. and Tally, P., 2007. Development of a health management strategy for the Silver Perch Aquaculture Industry. Final report on the Fisheries Research & Development Corporation, Project No. 2000/267 and 2004/089. NSW DPI – Fisheries Final Report Series No. 93. 219pp.
- No. 94 Park, T., 2007. NSW Gamefish Tournament Monitoring – Angling Research Monitoring Program. Final report to the NSW Recreational Fishing Trust. NSW DPI – Fisheries Final Report Series No. 94. 142pp.
- No. 95 Heasman, M.P., Liu, W., Goodsell, P.J., Hurwood D.A. and Allan, G.L., 2007. Development and delivery of technology for production, enhancement and aquaculture of blacklip abalone (*Haliotis rubra*) in New South Wales. Final Report to Fisheries Research and Development Corporation for Project No. 2001/33. NSW DPI – Fisheries Final Report Series No. 95. 226pp.
- No. 96 Ganassin, C. and Gibbs, P.J., 2007. A review of seagrass planting as a means of habitat compensation following loss of seagrass meadow. NSW Department of Primary Industries – Fisheries Final Report Series No. 96. 41pp.
- No. 97 Stewart, J. and Hughes, J., 2008. Determining appropriate harvest size at harvest for species shared by the commercial trap and recreational fisheries in New South Wales. Final Report to the Fisheries Research & Development Corporation for Project No. 2004/035. NSW Department of Primary Industries – Fisheries Final Report Series No. 97. 282pp.
- No. 98 West, G. and Williams, R.J., 2008. A preliminary assessment of the historical, current and future cover of seagrass in the estuary of the Parramatta River. SW Department of Primary Industries – Fisheries Final Report Series No. 98. 61pp.
- No. 99 Williams, D.L. and Scandol, J.P., 2008. Review of NSW recreational fishing tournament-based monitoring methods and datasets. NSW Department of Primary Industries – Fisheries Final Report Series No. 100. 83pp.
- No. 100 Allan, G.L., Heasman, H. and Bennison, S., 2008. Development of industrial-scale inland saline aquaculture: Coordination and communication of R&D in Australia. Final Report to the Fisheries Research & Development Corporation for Project No. 2004/241. NSW Department of Primary Industries – Fisheries Final Report Series No. 99. 245pp.
- No. 101 Gray, C.A and Barnes, L.M., 2008. Reproduction and growth of dusky flathead (*Platycephalus fuscus*) in NSW estuaries. NSW Department of Primary Industries – Fisheries Final Report Series No. 101. 26pp.