

## Grey Reef Shark, *Carcharhinus amblyrhynchos*

Report Card assessment	Sustainable		
IUCN Red List Australian Assessment	Near Threatened	IUCN Red List Global Assessment	Endangered
Assessors	Simpfendorfer, C., Fahmi, Bin Ali, A., , D., Utzurrum, J.A.T., Seyha, L., Maung, A., Bineesh, K.K., Yuneni, R.R., Sianipar, A., Haque, A.B., Tanay, D., Gautama, D.A. & Vo, V.Q.		
Australian Assessors	Kyne, P.M., Heupel, M.R., White, W.T. & Simpfendorfer, C.A. (Shark Action Plan)		
Report Card Remarks	Increasing numbers in the Great Barrier Reef Marine Park following rezoning and reduced Indonesian fishing pressure in the Timor Box.		

### Summary

The Grey Reef Shark is a common, large bodied species that inhabits coastal waters and oceanic atolls throughout the tropical Indo-Pacific. Site fidelity, restricted habitat, small litters, and late age at maturity have made it sensitive to increasing fishing pressure. Declines in local populations of over 90% have been recorded in the Indian Ocean and Western Central Pacific Ocean in recent decades. In Northern Australia,



Source: Will White. License: All rights reserved.

Indonesian fishermen considerably reduced the number of Grey Reef Sharks at Scott Reef in the 1990's. In the Great Barrier Reef Marine Park, there have been some declines in Grey Reef Shark but the levels of decline are uncertain. Rezoning of the Marine Park in 2004 appears to have promoted an increase in numbers of Grey Reef Sharks. Therefore, in Australia is assessed as Near Threatened (IUCN) (Kyne et al. 2021) and Sustainable (SAFS). From November 2023, the species will be listed on Appendix II of the Convention on International Trade in Endangered Species of Wild Flora and Fauna (CITES).

### Distribution

The Grey Reef Shark is a widespread, common species that occurs in the central Pacific and Indian Ocean (Last and Stevens 2009). It is found in Madagascar, Seychelles, and Reunion Island regions in the Indian Ocean. In the Pacific, it extends from southern China and Hawaii in the north to northern Australia in the south, and as far east as Easter Island. It is found in Northern Australia from south as Carnarvon (Western Australia), across the Northern Territory to Bundaberg (Queensland) and Lord Howe Island (Last and Stevens 2009).

### Stock structure and status

Within Australia, there is differing abundance levels of Grey Reef Shark depending on location. In northwest Australia at Scott Reef (within the Memorandum of Understanding [MOU] 1974 Box in the Timor Sea, where access by Indonesian fishers using traditional artisanal fishing techniques is permitted), the Indonesian fishermen caused substantial declines in Grey Reef Sharks throughout the 1990's to the point where Indonesian shark fishing vessels became uncommon (Wallner and McLoughlin 1996, Fox and Sen 2002, Meekan and Cappo 2004, Field et al. 2009). In the Great Barrier Reef Marine Park, there were significantly fewer Grey Reef Sharks in areas open to fishing and further declines in their population predicted (Robbins et al. 2006). A ten-year survey (2000–2010) that encompassed later years than the Robbins (et al. 2006) study found the numbers of Grey Reef Sharks increased following a major rezoning of the Marine Park in 2004, suggesting the zoning changes benefited the Grey Reef Shark (Espinoza et al. 2014). Globally, declines of over 90% have been recorded in the Indian and Western Central Pacific Oceans (White 2007, Graham et al. 2010, Nadon et al. 2012). It is estimated that the Grey Reef Shark has declined by 30% or more globally.

### Fisheries

Globally, it is targeted and taken as bycatch in artisanal and high seas longline, gillnet and trawl fisheries throughout its distribution (Kumoru 2003, White 2007, Graham et al. 2010). The Grey Reef Sharks are sensitive to fishing pressure due to high site fidelity, small litter size and late age at maturity. The species is retained for their fins and meat (Kumoru 2003). In Northern Australia, they have been fished heavily within the Timor Box and illegal, unregulated and unreported (IUU) fishing was a large and prevalent threat during the early 2000s (Field et al. 2009). However, the IUU fishing has declined significantly since 2006 due to increased border surveillance, international agreements and possibly increased fuel prices and the reduction in target species such as the Grey Reef Shark (Field et al. 2009, Haward and Bergin 2016). The Grey Reef Sharks have a strong affinity for sites with high coral cover (Espinoza et al. 2014), making them sensitive to habitat degradation. Even if fishing is well managed, threats affecting the quality of coral reef habitats may affect the population of Grey Reef Sharks. Within Australia, Grey Reef Sharks are taken in low numbers as bycatch in line fisheries targeting Coral Trout, however catch per unit effort data from 1986–2006 show no evidence of increase or decline of sharks taken in this fishery (Heupel et al. 2009).

### Habitat and biology

The Grey Reef Shark is found in tropical reef and continental shelf waters, occurring to depths of 280 m (Last and Stevens, 2009). It inhabits fringing reefs, lagoons and coral reef atolls selecting for areas with high coral cover and low turbidity (Wetherbee et al. 1997, Speed et al. 2011; Espinoza et al. 2014). The maximum recorded size is 190 cm total length (TL) and maximum age estimated is 19 years (Robbins 2006). Males mature at about 9 years and females at 11 years (Robbins 2006). Litters are small with mostly 3–4 pups (Robbins 2006).

Longevity and maximum size	Longevity: estimated 19 years Max size: 190 cm TL
Age and/or size at maturity (50%)	Males: 9 years, 120–140 cm TL Females: 11 years, 125–140 cm TL

**CAAB Code:** 37 018030

**Link to IUCN Page:** <https://www.iucnredlist.org/species/39365/173433550>

**Link to page at Shark References:** <http://shark-references.com/species/view/Carcharhinus-amblyrhynchos>

## References

- Espinoza, M., Cappelletti, M., Heupel, M.R., Tobin, A.J. and Simpfendorfer, C.A. 2014. Quantifying shark distribution patterns and species-habitat associations: implications of Marine Park Zoning. *PLOS ONE* 9(9): e106885.
- Field, I.C., Meekan, M.G., Buckworth, R.C. and Bradshaw, J.A. 2009. Protein mining the world's oceans. Australasia as an example of illegal expansion-and-displacement fishing. *Fish and Fisheries* 10: 323–328.
- Fox, J.J. and Sen, S. 2002. *A study of socio-economic issues of traditional Indonesian fishers who access the MOU Box*. Report for Environment Australia. Canberra.
- Graham, N.A.J., Spalding, M.D. and Sheppard, C.R.C. 2010. Reef shark declines in remote atolls highlight the need for multi-faceted conservation action. *Aquatic Conservation: Marine and Freshwater Ecosystems* 20: 543–548
- Haward, M. and Bergin, A. 2016. *Net worth: Australia's regional fisheries engagement*. Australian Strategic Policy Unit, Canberra.
- Heupel, M.R., Williams, A., Welch, D., Ballagh, A., Mapstone, B., Carlos, G., Davies, C. and Simpfendorfer, C.A. 2009. Effects of fishing on tropical reef associated shark populations on the Great Barrier Reef. *Fisheries Research* 95: 350–361.
- Kumoru, L. 2003. *The shark longline fishery in Papua New Guinea*. 176th Meeting of the Standing Committee on Tuna and Billfish: 5. Mooloolaba, Australia.
- Kyne, P.M., Heupel, M.R., White, W.T. and Simpfendorfer, C.A. 2021. *The Action Plan for Australian Sharks and Rays 2021*. National Environmental Science Program, Marine Biodiversity Hub, Hobart
- Last, P.R. and Stevens, J.D. 2009. *Sharks and Rays of Australia*. Second Edition. CSIRO Publishing, Collingwood, Australia.
- Meekan, M. and Cappelletti, M. 2004. *Non-destructive techniques for rapid assessment of shark abundance in Northern Australia*. The Australian Institute of Marine Science, Townsville.
- Nadon, Marc O; Baum, Julia K; Williams, Ivor D; Mcpherson, Jana M; Zgliczynski, Brian J; Richards, Benjamin L; Schroeder, Robert E; Brainard, Russell E. 2012. Re-creating missing population baselines for Pacific reef sharks. *Conservation Biology* 26(3): 493–503.
- Robbins, W.D., Hisano, M., Connolly, S.R. and Choat, J.H. 2006. Ongoing collapse of coral-reef shark populations. *Current Biology* 16: 2314–2319.
- Speed, Conrad W; Meekan, Mark G; Field, Iain C; McMahon, Clive R; Stevens, John D; McGregor, Frazer; Huvener, Charlie; Berger, Yuval; Bradshaw, Corey J A. 2011. Spatial and temporal movement patterns of a multi-species coastal reef shark aggregation. *Marine Ecology Progressive Series* 429: 261–275.
- Wallner, B.K. and McLoughlin, K. 1996. Review of Indonesian fishing in the Australian fishing zone. Report for Fisheries Policy Branch. Department of Primary Industries and Energy, Canberra.
- Wetherbee, B.M., Crow, G.L. and Lowe, C.G. 1997. Distribution, reproduction and diet of the gray reef shark, *Carcharhinus amblyrhynchos* in Hawaii. *Marine Ecology Progress Series* 151: 181–189.
- White, W.T. 2007. Catch composition and reproductive biology of whaler sharks (Carcharhiniformes: Carcharhinidae) caught by fisheries in Indonesia. *Journal of Fish Biology* 71(5): 1510–1540.