

Silky Shark, *Carcharhinus falciformis*

Report Card assessment	Depleting		
IUCN Red List Australian Assessment	Vulnerable	IUCN Red List Global Assessment	Vulnerable
Global Assessors	Rigby, C.L., Sherman, C.S., Chin, A. & Simpfendorfer, C.		
Australian Assessors	Kyne, P.M., Heupel, M.R., White, W.T. & Simpfendorfer, C.A. (Shark Action Plan)		
Report Card Remarks	In Australia, there is no take of Silky Sharks in the tuna fisheries. Listed on CITES Appendix II, CMS Appendix II.		

Summary

The Silky Shark is a large bodied oceanic and coastal-pelagic shark that has a global distribution throughout tropical waters. It is taken as bycatch or as a target species, and is the second most caught pelagic shark species globally after Blue Sharks (*Prionace glauca*). Silky Sharks have a long generation length of 15 years (average age of parents) and are thus sensitive to fishing pressure. In some regions the catches have remained stable or slightly increased,



Source: Joi Ito/Flickr. License: CC By Attribution.

while in others declines of up to 99% have been estimated. There is a very high level of uncertainty in the estimates of decline and an inability to conclusively attribute declines solely to fishing mortality as there is some potential for environmental influences on catchability. The weighted population global decline over three generations is estimated to be 47–54%. Therefore, the Silky Shark is assessed globally as Vulnerable (IUCN); and in Australia as Vulnerable (IUCN) as it is part of shared regional stocks (Kyne et al. 2021), and Depleting (SAFS). The species is listed on CITES Appendix II and CMS Appendix II.

Distribution

The Silky Shark is globally distributed throughout tropical pelagic waters and may move seasonally into warm temperate waters (Last and Stevens 2009). Within Australia, it is found from Lancelin (Western Australia), through the Northern Territory and Queensland to Sydney (New South Wales) (Last and Stevens 2009). Globally, it is found throughout the tropics across the Atlantic, Indian and Pacific Oceans.

Stock structure and status

There is no information on the stock structure in Australian waters. However, animals caught in Australian waters are part of regional stocks in the Indian and Pacific oceans. Within Australia, there is currently no information on population size, or trend for the Silky Shark. Life history parameters vary geographically, suggesting distinct populations. Within the Pacific, there are potentially three stocks; one in the western Pacific and two in the eastern Pacific that are separated by the equator (Aires da Silva et al. 2014). In the western Pacific, the population exhibited no strong trend of decline over 1995–2014 (Rice et al. 2015). In the eastern Pacific, declines in the north ranged from 17–60% over three generations due to uncertainties in the catch data and in the south more definitive declines of up to 99% are reported (Lennart-Cody 2015). In the Atlantic, declines are also reported of up to 95% over three generations (Cortes et al. 2007). When the three generation population trend is weighted according to the relative size of each region, the global population decline is estimated to be 47–54% (Rigby et al. 2021). Given that animals caught in Australian waters are part of regional stocks in the Indian and Western Pacific oceans that are declining, but that may not yet be overfished, the species meets SAFS criteria as Depleting.

Fisheries

Silky Sharks are retained for their meat and fins and in some fisheries, such as the tropical purse seine fisheries, their mortality remains high even when released (Hutchinson et al. 2015). Silky Sharks are sensitive to fishing pressure due to their life history characteristics. Within Australia, the catches are likely minimal in the west where pelagic longlining fishing effort is very low. There are now bans on the take of Silky Sharks in both the Eastern and Western Tuna and Billfish Fisheries (AFMA 2015, 2016) and so any caught are released. Globally, fishing pressure is high with Silky Sharks the second most caught species of shark (Oliver et al. 2015).

Habitat and biology

The Silky Shark is found in pelagic and continental shelf edge waters throughout the tropics and occurs from the surface to depths of 500 m (Last and Stevens 2009). Maximum age and size range vary geographically. Maximum observed size is 330 cm total length (TL) and maximum estimated size is approximately 370 cm TL (Serafy et al. 2012, Ebert et al. 2013). Maximum age estimates vary from 16 to 36 years (Sanchez-de-Ita et al. 2011, Joung et al. 2008).

Longevity and maximum size	Longevity: estimates from 16 to 36 years Max size: 330 cm TL (observed)
Age and/or size at maturity (50%)	Males: 180–225 cm TL Females: 180–246 cm TL

CAAB Code: 37 018008

Link to IUCN Page: <https://www.iucnredlist.org/species/39370/205782570>

Link to page at Shark References: <http://shark-references.com/species/view/Carcharhinus-falciformis>

References

- Aires-da-Silva, A., Lennert-Cody, C., Maunder, M.N. and Román-Verdesoto, M. 2014. *Stock status indicators for silky sharks in the eastern Pacific Ocean*. Document SAC-05-11a. Inter-American Tropical Tuna Commission Scientific Advisory Committee Fifth Meeting. 12-16 May 2014, La Jolla, California, USA [<http://www.iattc.org/Meetings/Meetings2014/MAYSAC/PDFs/SAC-05-11a-Indicators-for-silky-sharks.pdf>].
- Australian Fisheries Management Authority. 2015. *Eastern Tuna and Billfish Fishery (ETBF). Management arrangements booklet*. 2015 Fishing Season.
- Australian Fisheries Management Authority. 2016. *Western Tuna and Billfish Fishery (WTBF). Management arrangements booklet*. 2016 Fishing Season.

- Cortés, E., Brown, C. and Beerkircher, L.R. 2007. Relative abundance of pelagic sharks in the Western North Atlantic Ocean, including the Gulf of Mexico and Caribbean Sea. *Gulf and Caribbean Research* 19(2): 37–52.
- Ebert, D.A., Fowler, S. and Compagno, L. 2013. *Sharks of the world. A fully illustrated guide*. Wild Nature Press.
- Hutchinson, M.R., Itano, D.G., Muir, J.A. and Holland, K.N. 2015. Post-release survival of juvenile silky sharks captured in a tropical tuna purse seine fishery. *Marine Ecology Progress Series* 521: 143–154.
- Joung, S.J., Chen, C., Lee, H. and Liu, K. 2008. Age, growth and reproduction of silky sharks, *Carcharhinus falciformis*, in northeastern Taiwan waters. *Fisheries Research*, 90, 78–85
- Kyne, P.M., Heupel, M.R., White, W.T. and Simpfendorfer, C.A. 2021. *The Action Plan for Australian Sharks and Rays 2021*. National Environmental Science Program, Marine Biodiversity Hub, Hobart
- Last, P.R. and Stevens, J.D. 2009. *Sharks and Rays of Australia*. Second Edition. CSIRO Publishing, Collingwood.
- Lennert-Cody, C., Aires-da-Silva, A., Maunder, M. and Roman, M. H. 2016. *Updated stock status indicators for Silky Sharks in the eastern Pacific Ocean*. Inter-American Tropical Tuna Commission Scientific Advisory Committee Seventh Meeting, La Jolla, California.
- Oliver, S., Braccini, M., Newman, S.J. and Harvey, E.S. 2015. Global patterns in the bycatch of sharks and rays. *Marine Policy* 54: 86–97.
- Rice, J., Tremblay-Boyer, L., Scott, R., Hare, S. and Tidd, A. 2015. *Analysis of stock status and related indicators for key shark species of the Western Central Pacific Fisheries Commission*. In: Western Central Pacific Fisheries Commission (ed.), Scientific Committee Eleventh Regular Session, 5-13 August 2015, Pohnpei, Federated States of Micronesia.
- Rigby, C.L., Sherman, C.S., Chin, A. & Simpfendorfer, C. 2021. *Carcharhinus falciformis* (amended version of 2017 assessment). *The IUCN Red List of Threatened Species* 2021: e.T39370A205782570.
- Sánchez-de Ita, J.A., Quiñónez-Velázquez, C., Galván-Magaña, F., Bocanegra-Castillo, N. and Félix-Uraga, R. 2011. Age and growth of the silky shark *Carcharhinus falciformis* from the west coast of Baja California Sur, Mexico. *Journal of Applied Ichthyology* 27: 20–24.
- Serafy, J.E., Orbesen, E.S., Snodgrass, D.J., Beerkircher, L.R. and Walter, J.F. 2012. Hooking survival of fishes captured by the United States Atlantic pelagic longline fishery: impact of the 2004 circle hook rule. *Bulletin of Marine Science* 88(3): 605–621.