

## Bigeye Sixgill Shark, *Hexanchus nakamurai*

Report Card assessment	Negligible		
IUCN Red List Australian Assessment	Least Concern	IUCN Red List Global Assessment	Near Threatened
Assessors	Finucci, B., Barnett, A., Cheok, J., Cotton, C.F., Kulka, D.W., Neat, F.C., Rigby, C.L., Tanaka, S. & Walker, T.I.		
Report Card Remarks	Rarely encountered in Australia fisheries.		

### Summary

The Bigeye Sixgill Shark is a little known, mainly deepwater species with a wide ranging but patchy distribution. It is uncommonly taken by trawl and longline gear and is of relatively minor importance to fisheries. The species has often been misidentified as the larger Bluntnose Sixgill Shark (*Hexanchus griseus*),



leading to confusion and poor knowledge of its distribution, catches and population trends. In Australia, it is rarely encountered in fisheries. Therefore, the species is assessed as Least Concern (IUCN) in Australia (Kyne et al. 2021) and Negligible (SAFS) due to its rare interaction with fisheries.

### Distribution

The Bigeye Sixgill Shark occurs across tropical and warm-temperate waters of the western Pacific, Atlantic and Indian Oceans, but appears to be absent from the entire eastern Pacific Ocean. In Australia, it is recorded from Queensland down to Tathra (New South Wales) and off the North West Shelf and Bunbury (Western Australia) (Last and Stevens, 2009).

### Stock structure and status

There is currently no information on population size, structure, or trend for the species. It is uncommon to rare where it occurs (Ebert 1990, Serena 2005).

### Fisheries

The species is taken as bycatch on line and trawl fisheries. In southeast Australia, where it is rarely caught in the Southern and Eastern Scalefish and Shark Fishery, the species is considered rare based on reported catches. It has a moderate to low catch susceptibility to trawl, gillnet, and hook fishing gears and is categorized as at high ecological risk (Walker et al. 2008). The species has also been recorded in very low numbers in catches of the artisanal deepwater longline fishery operating off eastern Indonesia, where all landed elasmobranchs are utilised (White and Dharmadi, 2010). Small

numbers are caught off Campeche Bank, Yucatán in Mexican shark fisheries, as bycatch of snapper fisheries in the Cayman Islands, and as bycatch of trawl fisheries off Taiwan (Ebert 1990).

### Habitat and biology

The Bigeye Sixgill Shark is found on the continental and insular shelves and slopes from 60–620 m depth, usually on or near bottom, but occasionally moving to near the surface or inshore in the tropics (Ebert 1990, Barnett et al. 2012). Maximum size is approximately 180 cm total length (TL). Males mature at 142–178 cm TL and females at 123–157 cm TL (Ebert 1990, Barnett et al. 2012).

Longevity and maximum size	Longevity: unknown Max size: 180 cm TL
Age and/or size at maturity (50%)	Males: 142–178 cm TL Females: 123–157 cm TL

**CAAB Code:** 37 005004

**Link to IUCN Page:** <https://www.iucnredlist.org/species/152781341/68626047>

**Link to page at Shark References:** <http://www.shark-references.com/species/view/Hexanchus-nakamurai>

#### References

- Barnett, A. Braccini, J.M. Awruch, C. A. Ebert, D.A. 2012. An overview on the role of Hexanchiformes in marine ecosystems: biology, ecology and conservation status of a primitive order of modern shark. *Journal of Fish Biology* 80: 966–990.
- Ebert, D.A. 1990. *The taxonomy, biogeography and biology of cow and frilled sharks (Chondrichthyes: Hexanchiformes)*. Unpub. Ph.D. thesis. Rhodes University.
- Kyne, P.M., Heupel, M.R., White, W.T. and Simpfendorfer, C.A. 2021. *The Action Plan for Australian Sharks and Rays 2021*. National Environmental Science Program, Marine Biodiversity Hub, Hobart
- Last P.R. and Stevens J.D. 2009. *Sharks and Rays of Australia*, Second Edition. CSIRO Publishing, Melbourne.
- Serena, F. 2005. *Field identification guide to the sharks and rays of the Mediterranean and Black Sea*.
- Walker T.I., Stevens J.D., Braccini J.M., Daley R.K., Huveneers C., Irvine SB., Bell J.D., Tovar-Avila J., Trinnie F.I., Phillips D.T., Treloar M.A., Awruch C.A., Gason A.S., Salini J. & Hamlett W.C. 2008. *Rapid Assessment of Sustainability for Ecological Risk of Shark and Other Chondrichthyan Bycatch Species Taken in the Southern and Eastern Scalefish and Shark Fishery*. Primary Industries Research Victoria, Queenscliff.
- White W.T. and Dharmadi. 2010. Aspects of maturation and reproduction in hexanchiform and squaliform sharks. *Journal of Fish Biology* 76: 1362–1378.