

Tiger Shark, *Galeocerdo cuvier*

Report Card assessment	Depleting		
IUCN Red List Australian Assessment	Refer to Global Assessment	IUCN Red List Global Assessment	Near Threatened
Assessors	Ferreira, L.C. & Simpfendorfer, C.		
Report Card Remarks	Declines in catch rates in Australian Shark Control Programs		

Summary

The Tiger Shark is a large, highly mobile shark species globally distributed throughout tropical to temperate waters. It is targeted and taken as bycatch by commercial, recreational and artisanal fisheries throughout much of its distribution. It is thought to reproduce only every three years which reduces its



Source: Kevin Bryant/Flickr. License: CC By Attribution-NonCommercial-ShareAlike.

capacity to recover from exploitation. Declines in Tiger Shark catch rates have been recorded in the Queensland Shark Control Program and in the New South Wales Shark Meshing Program. Increases in catch rates have been recorded in South Africa beach protection program (from 1978-2003) and both coasts of the United States, however these increases are minimal and should be interpreted with caution. Little information exists on catches throughout the rest of its distribution. Catch rate trends and information on population structure from shark control programs in Australia should continue to be monitored due to concerns over the effect those programs are having on the Tiger Shark population off eastern Australia. Off western Australia, catches of Tiger Sharks are low in the commercial fisheries. Therefore, the Tiger Shark is assessed as Near Threatened (IUCN) and in Australia, Transitional Depleting (SAFS) due to concerns of catches in the shark control programs.

Distribution

The Tiger Shark has a circumglobal distribution throughout tropical and temperate waters (Compagno 1984). Within Australia, it is found from Windy Harbour (Western Australia) through the Northern Territory, Queensland and to Sydney (New South Wales) (Last and Stevens 2009).

Stock structure and status

There is currently limited information on population size, structure, or trend for Tiger Sharks. It is not captured in large numbers by most commercial fisheries compared to other pelagic sharks and the catch is rarely recorded. Within Australia, in the Queensland Shark Control Program, the catch rate increased from 1962 until the 1980s and then decreased in the 2000s to approximately 66% of its original value; the most significant declines were in the southern region of the State (Simpfendorfer et al. 2010, Holmes et al. 2012). In New South Wales (NSW), the catch rate and size of Tiger Sharks has

also declined in the last two decades which raises concerns over the effect the program is having on the Tiger Shark population off NSW (Reid et al. 2011). In the Northwest Atlantic, increases in relative abundance have been recorded from 1960-2000 by the Northeast Fisheries Science Centre and no change in abundance recorded by the Southeast Fisheries Science Centre for the same period (J. Carlson, pers. comm). The KwaZulu Natal beach protection program showed a rate of change of 3% increase per year in Tiger Shark catch between 1978 and 2003. The increasing rates in the Northwest Atlantic and South Africa seem consistent with the pattern observed in the first 20 years in Queensland and the interpretation of such upward trends should be made with caution. With the limited data available indicates that the significant drivers of population decline at present are the shark control programs in Australia and South Africa. Given the wide range of this species and the localities of the programs, the Tiger Shark is suspected to have undergone an overall population reduction of <30% globally.

Fisheries

The Tiger Shark is targeted and taken as bycatch in a number of commercial, artisanal and recreational fisheries throughout its distribution for its meat and fins. Fishing pressure has been gradually increasing since the 1950's due to the demand from the shark fin trade (Ward-Paige et al. 2010). Within Australia, Tiger Sharks are taken incidentally in the Offshore Net and Line Fishery in northern Australia (27 t in 2012; DPFI 2012). The species is taken in the Southern and Western Demersal Gillnet and Demersal Longline Fishery in Western Australia with a catch of 112 t reported in 2005-2006 (McAuley 2008). In northwest Australian waters, shark fishing gear has been prohibited in those fisheries since 1993 and further north, both the West Australian Northern Shark Fishery and the Joint Authority Northern Shark Fishery have not operated since 2009 (Fletcher and Santoro 2013). Prior to that varying catches of 12-81 t of Tiger Shark were taken annually from 2001-2002 to 2005-2006 from those two fisheries (Fletcher and Santoro 2008). The Australian Commonwealth Trawl Sector reported a total catch for the species of 4.7 t from 2004-2011. In the game fishery off the southeast coast of Australia, the species comprised 10% of catches from 1961-1990 with individuals ranging in mass from 21-560 kg (Pepperell 1992). Tiger Shark catches in the New South Wales Gamefish Tournament were approximately 8 t per year, comprising approximately 1-5% of the total catch and 10-30% of shark catches between 1993 and 2005 (Park 2007). The Tiger Shark is targeted by Shark Control Programs in Australia and South Africa (Dudley and Simpfendorfer 2006, Cliff and Dudley 2011, Holmes et al. 2012). The Queensland Shark Control Program captured a total of 4,757 individuals from 1993-2010 (Holmes et al. 2012). The New South Wales Shark Meshing Program captured approximately 30 sharks per year from 1950-2008; an approximate total of 1,740 individuals.

Habitat and ecology

The Tiger Shark is encountered throughout tropical and temperate waters on coral reefs, continental shelves and offshore islands and atolls. It occurs across a range of depths, from very shallow waters to depths of 1,136m (Heithaus et al. 2002, Werry et al. 2014). The maximum age is 27-33 years old (Branstetter et al. 1987, Wintner and Dudley 2000, Kneebone et al. 2008). Estimates of size and age at maturity vary greatly between regions. Although it has large litters of 3-82 pups, it is estimated to reproduce only once every three years (Whitney and Crow 2007, B. Holmes pers. comm.).

Longevity and maximum size	Longevity: estimated 27-33 years Max size: at least 600 cm TL
Age and/or size at maturity (50%)	Both sexes: 5-13 years Males: ~250-305 cm TL Females: 274-345 cm TL

Link to IUCN Page: <https://www.iucnredlist.org/species/39378/10220026>

Link to page at Shark References: <http://shark-references.com/species/view/Galeocerdo-cuvier>

References

- Branstetter, S., Musick, J.A. and Colvocoresses, J.A. 1987. A comparison of the age and growth of the tiger shark, *Galeocerdo cuvieri*, from off Virginia and from the northwestern Gulf of Mexico. *NMFS Fishery Bulletin* 85: 269–79.
- Cliff, G., Dudley, S.F.J. 2011. Reducing the environmental impact of shark-control programs: a case study from KwaZulu-Natal, South Africa. *Marine and Freshwater Research* 62: 700-709.
- Compagno, L.J.V. 1984. FAO species catalogue. Vol. 4. Sharks of the world. An annotated and illustrated catalogue of shark species known to date. FAO Fish. Synop. No. 125, vol. 4.
- DPFI 2012. Offshore Net and Line Fishery Status Report 2012. Fishery Report No 113. Department Fisheries.
- Dudley, S.F.J. and Simpfendorfer, C.A. 2006. Population status of 14 shark species caught in the protective gillnets off KwaZulu-Natal beaches, South Africa, 1978-2003. *Marine and Freshwater Research* 57: 225-240.
- Fletcher, W.J. and Santoro, K. (eds). 2008. State of the Fisheries Report 2007/08. Department of Fisheries, Western Australia.
- Fletcher, W.J. and Santoro, K. (eds). 2013. Status Reports of the Fisheries and Aquatic Resources of Western Australia 2012/13: The State of the Fisheries. Department of Fisheries, Western Australia.
- Heithaus, M.R., Dill, L.M., Marshall, G.J., Buhleier, B. 2002. Habitat use and foraging behavior of tiger sharks (*Galeocerdo cuvier*) in a seagrass ecosystem. *Marine Biology* 140: 237-248.
- Holmes, B.J., Sumpton, W.D., Mayer, D.G., Tibbetts, I.R., Neil, D.T., Bennett, M.B. 2012. Declining trends in annual catch rates of the tiger shark (*Galeocerdo cuvier*) in Queensland, Australia. *Fisheries Research* 129-130: 38-45.
- Kneebone, J., Natanson, L.J., Andrews, A.H., Howell, W.H. 2008. Using bomb radiocarbon analyses to validate age and growth estimates for the tiger shark, *Galeocerdo cuvier*, in the western North Atlantic. *Marine Biology* 60: 1-87.
- Last, P.R. and Stevens, J.D. 2009. Sharks and Rays of Australia. Second Edition. CSIRO Publishing, Collingwood, Australia.
- McAuley, R. 2008. Demersal Gillnet and Longline Fisheries Status Report. State of Fisheries Report 2006/2007. Department of Fisheries, Perth.
- Park, T. 2007. NSW Gamefish Tournament Monitoring – Angling Research Tournament Monitoring Program. NSW Department of Primary Industries, Cronulla, NSW
- Pepperell, J. 1992. Trends in the distribution, species composition and size of sharks caught by gamefish anglers off south-eastern Australia, 1961?1990. In: J. Pepperell (ed.), Sharks: Biology and Fisheries. *Australian Journal of Marine and Freshwater Research*, pp. 213–225.
- Reid, D.D., Robbins, W.D., Peddemors, V.M. 2011. Decadal trends in shark catches and effort from the New South Wales, Australia, Shark Meshing Program 1950-2010. *Marine and Freshwater Research* 62: 676-693.
- Simpfendorfer, C.A., de Jong, S.K., Sumpton, W. 2010. Long-term trends in large shark populations from inshore areas of the Great Barrier Reef World Heritage Area: results from the Queensland Shark Control Program. Marine and Tropical Sciences Research Facility Transition Program Report.
- Ward-Paige, C.A., Mora, C., Lotze, H.K., Pattengill-Semmens, C., McClenachan, L., Arias-Castro, E., Myers, R.A. 2010. Large-scale absence of sharks on reefs in the greater-Caribbean: a footprint of human pressures. *Plos One* 5: e11968.
- Werry, J.M., Planes, S., Berumen, M.L., Lee, K.A., Braun, C.D., Clua, E. 2014. Reef Fidelity and Migration of Tiger Sharks, *Galeocerdo cuvier*, across the Coral Sea. *Plos One* 9: e83249.
- Whitney, N.M., Crow, G.L. 2007. Reproductive biology of the tiger shark (*Galeocerdo cuvier*) in Hawaii. *Marine Biology* 151: 63-70.
- Wintner, S.P., Dudley, S.F.J. 2000. Age and growth estimated for the tiger shark, *Galeocerdo cuvier*, from the east coast of South Africa. *Marine and Freshwater Research* 51: 43-53.